Author Index

Agostini, O.
—, Moneti, G., Bonacchi, G., Fedi, M. and Manzini, S.
Determination of the antihypertensive drug 1-[2-ethoxy-2-(3′-pyridyl)ethyl]-4-(2′-methoxyphenyl)piperazine (IP/66) in rat and human plasma by high-performance liquid chromatography and isotope dilution mass spectrometry 331
d'Athis, P., see Tsaconas, C. 313
Aubert, C., see Drouet-Coassolo, C. 295
Benny, A.G.
—, Ockelford, P.A., Brimer, C.M. and Morgan, A.K.
Separation of vitamin K-dependent coagulation factors on sulphated cellulose 173
Blaauboer, B.J., see Van Pelt, F.N.A.M. 489
Bleau, G.
— and Desaulniers, M.
High-performance liquid chromatographic assay of benzalkonium in plasma 221
Bonacchi, G., see Agostini, O. 331
Boomsma, F., see Van der Hoorn, F.A.J. 17
Boppana, V.K.
—, Dolce, K.M., Cyronak, M.J. and Ziemniak, J.A.
Simplified procedures for the determination of fenoldopam and its metabolites in human plasma by high-performance liquid chromatography with electrochemical detection: comparison of manual and robotic sample preparation methods 385
Boutagy, J.
—, Rumble, F. and Dunagan, F.
Determination of isradipine and the oxidative pyridine metabolite in human plasma by high-performance liquid chromatography 483
Bowers, L., see Edlund, P.O. 341
Brimer, C.M., see Benny, A.G. 173
Brittain, H.G., see Kadin, H. 135
Bromet, N., see Tsaconas, C. 313
Brooks, B.R., see Turner, J.G. 275
Brouwer, K.L.R., see Jewell, R.C. 257
Brown, N.D.
—, Poon, B.T. and Phillips, L.R.
Identification and determination of a carboxylic acid metabolite of chloroquine in human urine by high-performance liquid chromatography 189
Bryan, L.J.
— and O'Donnell, S.R.
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by high-performance liquid chromatography with electrochemical detection 29
Bryn, K., see Sonesson, A. 1
Burger, J.A., see Labadarios, D. 428
Burstein, E.S.
— and Greenblatt, D.J.
Simplified gas chromatographic analysis of ethanol in blood and tissue 228
Cano, J.-P., see Drouet-Coassolo, C. 295
Capdepuy, M., see Fourche, J. 142
Case, G.L.
— and Steele, R.D.
Determination of reduced folate derivatives in tissue samples by high-performance liquid chromatography with fluorimetric detection 456
Chong, W.K.
—, Mills, G.A., Weavind, G.P. and Walker, V.
High-performance liquid chromatographic method for the rapid profiling of plasma and urinary organic acids 147
Clavey, V., see Steinmetz, A. 154
Coassolo, P., see Drouet-Coassolo, C. 295
Cohen, A.I., see Kadin, H. 135
Cyronak, M.J., see Boppana, V.K. 385
Czupryna, M.J., see Saller, C.F. 167
Daas, I. den, see Den Daas, I. 210
Davis, P.F., see Manning, J.N. 41
Den Daas, I.
—, Rollema, H., De Vries, J.B., Tepper, P.G. and Horn, A.S.
Analysis of the dopamine agonist N-0437 in rat serum using reversed-phase high-performance liquid chromatography with electrochemical detection 210
Desaulniers, M., see Bleau, G. 221
Dessalles, M.C.
—, Vienne, B. and Mahuzier, G.
Dosage sérique de la perphénazine en chromatographie liquide 215
De Vries, J.B., see Den Daas, I. 210
De Zeeuw, R.A., see Gerding, T.K. 125
Dolce, K.M., see Boppana, V.K. 385
Downer, R.G.H., see Martin, R.J. 287
Drenth, B.F.H., see Gerding, T.K. 125

Author Index

Agostini, O.
—, Moneti, G., Bonacchi, G., Fedi, M. and Manzini, S.
Determination of the antihypertensive drug 1-[2-ethoxy-2-(3′-pyridyl)ethyl]-4-(2′-methoxyphenyl)piperazine (IP/66) in rat and human plasma by high-performance liquid chromatography and isotope dilution mass spectrometry 331
d'Athis, P., see Tsaconas, C. 313
Aubert, C., see Drouet-Coassolo, C. 295
Benny, A.G.
—, Ockelford, P.A., Brimer, C.M. and Morgan, A.K.
Separation of vitamin K-dependent coagulation factors on sulphated cellulose 173
Blaauboer, B.J., see Van Pelt, F.N.A.M. 489
Bleau, G.
— and Desaulniers, M.
High-performance liquid chromatographic assay of benzalkonium in plasma 221
Bonacchi, G., see Agostini, O. 331
Boomsma, F., see Van der Hoorn, F.A.J. 17
Boppana, V.K.
—, Dolce, K.M., Cyronak, M.J. and Ziemniak, J.A.
Simplified procedures for the determination of fenoldopam and its metabolites in human plasma by high-performance liquid chromatography with electrochemical detection: comparison of manual and robotic sample preparation methods 385
Boutagy, J.
—, Rumble, F. and Dunagan, F.
Determination of isradipine and the oxidative pyridine metabolite in human plasma by high-performance liquid chromatography 483
Bowers, L., see Edlund, P.O. 341
Brimer, C.M., see Benny, A.G. 173
Brittain, H.G., see Kadin, H. 135
Bromet, N., see Tsaconas, C. 313
Brooks, B.R., see Turner, J.G. 275
Brouwer, K.L.R., see Jewell, R.C. 257
Brown, N.D.
—, Poon, B.T. and Phillips, L.R.
Identification and determination of a carboxylic acid metabolite of chloroquine in human urine by high-performance liquid chromatography 189
Bryan, L.J.
— and O'Donnell, S.R.
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by high-performance liquid chromatography with electrochemical detection 29
Bryn, K., see Sonesson, A. 1
Burger, J.A., see Labadarios, D. 428
Burstein, E.S.
— and Greenblatt, D.J.
Simplified gas chromatographic analysis of ethanol in blood and tissue 228
Cano, J.-P., see Drouet-Coassolo, C. 295
Capdepuy, M., see Fourche, J. 142
Case, G.L.
— and Steele, R.D.
Determination of reduced folate derivatives in tissue samples by high-performance liquid chromatography with fluorimetric detection 456
Chong, W.K.
—, Mills, G.A., Weavind, G.P. and Walker, V.
High-performance liquid chromatographic method for the rapid profiling of plasma and urinary organic acids 147
Clavey, V., see Steinmetz, A. 154
Coassolo, P., see Drouet-Coassolo, C. 295
Cohen, A.I., see Kadin, H. 135
Cyronak, M.J., see Boppana, V.K. 385
Czupryna, M.J., see Saller, C.F. 167
Daas, I. den, see Den Daas, I. 210
Davis, P.F., see Manning, J.N. 41
Den Daas, I.
—, Rollema, H., De Vries, J.B., Tepper, P.G. and Horn, A.S.
Analysis of the dopamine agonist N-0437 in rat serum using reversed-phase high-performance liquid chromatography with electrochemical detection 210
Desaulniers, M., see Bleau, G. 221
Dessalles, M.C.
—, Vienne, B. and Mahuzier, G.
Dosage sérique de la perphénazine en chromatographie liquide 215
De Vries, J.B., see Den Daas, I. 210
De Zeeuw, R.A., see Gerding, T.K. 125
Dolce, K.M., see Boppana, V.K. 385
Downer, R.G.H., see Martin, R.J. 287
Drenth, B.F.H., see Gerding, T.K. 125

Drouet-Coassolo, C.
—, Aubert, C., Coassolo, P. and Cano, J.-P.
Capillary gas chromatographic–mass spectrometric method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295
Dubois, J.P., see Sioufi, A. 81
Dunagan, F., see Boutagy, J. 483
Edlund, P.O.
—, Bowers, L. and Henion, J.
Determination of methandrostenolone and its metabolites in equine plasma and urine by coupled-column liquid chromatography with ultraviolet detection and confirmation by tandem mass spectrometry 341
Everett, D.W.
—, Foley, J.E., Singhvi, S.M., Weinstein, S.H. and Warrington, S.J.
High-performance liquid chromatographic method for the radiometric determination of [^{14}C]bucromarone in human plasma utilizing non-radiolabeled bucromarone as an internal standard 365
Faulkner, R., see Nuernberg, B. 476
Fedi, M., see Agostini, O. 331
Fedorchuk, M., see Schmitthenner, H.F. 197
Foley, J.E., see Everett, D.W. 365
Fourche, J.
—, Capdepuy, M. and Texier-Maugein, J.
Gas chromatographic fatty acid determination to differentiate *Nocardia asteroides*, *Mycobacterium fortuitum* and *Mycobacterium chelonei* 142
Fruchart, J.-C., see Steinmetz, A. 154
Fu, C.-H.J., see Gillilan, R.B. 232
Furst, D.E., see Nuernberg, B. 476
Gerding, T.K.
—, Drenth, B.F.H., Van de Grampel, V.J.M., Niemeijer, N.R., De Zeeuw, R.A., Tepper, P.G. and Horn, A.S.
Determination of enantiomeric purity of the new D-2 dopamine agonist 2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin (N-0437) by reversed-phase high-performance liquid chromatography after pre-column derivatization with D(+)-glucuronic acid 125
Gillilan, R.B.
—, Mason, W.D. and Fu, C.-H.J.
Rapid analysis of piroxicam in dog, rat and human plasma by high-performance liquid chromatography 232
Girometta, M.A.
—, Loschi, L. and Ventura, P.
High-performance liquid chromatographic assay of nesosteine in human plasma and urine 117
Gonzalez, M.A., see Ruggirello, D.A. 73
Grampel, V.J.M. van de, see Gerding, T.K. 125
Greenblatt, D.J., see Burstein, E.S. 228
Gu, L.H., see Kutlar, F. 265
Hassing, I.G.A.M., see Van Pelt, F.N.A.M. 489
Hawes, E.M., see Srinivas, N.R. 61
Hayakawa, K.
— and Oizumi, J.
Isocratic separation of phenylthiohydantoin-amino acids by reversed-phase high-performance liquid chromatography 161
Hendrickson, H.P.
—, Scott, D.O. and Lunte, C.E.
Identification of 9-hydroxylamine-1,2,3,4-tetrahydroacridine as a hepatic microsomal metabolite of tacrine by high-performance liquid chromatography and electrochemistry 401
Henion, J., see Edlund, P.O. 341
Ho, P., see Ruggirello, P. 73
Honegger, C.G.
—, Langemann, H., Krenger, W. and Kempf, A.
Liquid chromatographic determination of common water-soluble antioxidants in biological samples 463
Hoorn, F.A.J. van der, see Van der Hoorn, F.A.J. 17
Horn, A.S., see Den Daas, I. 210
—, see Gerding, T.K. 125
Hosotsubo, H.
Rapid and sensitive method for the determination of 5-fluorocytosine in human plasma by ion-pair high-performance liquid chromatography 183
Rapid and specific method for the determination of vancomycin in plasma by high-performance liquid chromatography on an aminopropyl column 421
—, Takeda, K., Hosotsubo, K. and Yoshiya, I.
Measurement of thiamylal in human plasma using reversed-phase high-performance liquid chromatography 204
Hotsubo, K., see Hosotsubo, H. 204
Hough, B.J., see Labadarios, D. 428
Hu, H., see Kutlar, F. 265
Hubbard, J.W., see Srinivas, N.R. 61
Huisman, T.H.J., see Kutlar, F. 265
Ishii, T., see Sugata, R. 9

Ivashkiv, E., see Kadin, H. 135
Iwahashi, H., see Sugata, R. 9
Jantzen, E., see Sonesson, A. 1
Jarvi, E.J., see Patrick, K.S. 91
Jewell, R.C.
—, Brouwer, K.L.R. and McNamara, P.J.
α_1-Acid glycoprotein high-performance liquid chromatography column (EnantioPAC) as a screening tool for protein binding 257
Kadin, H.
—, Brittain, H.G., Ivashkiv, E. and Cohen, A.I.
Body fluid analysis of a phosphonic acid angiotensin-converting enzyme inhibitor using high-performance liquid chromatography and post-column derivatization with *o*-phthalaldehyde 135
Kaffarnik, H., see Steinmetz, A. 154
Kaplan, B.
— and Pras, M.
Separation of peptides from sodium dodecyl sulphate by high-performance liquid chromatography 450
Kempf, A., see Honegger, C.G. 463
Kido, R., see Sugata, R. 9
Köhl, U., see Schwabe, D. 177
Kohlbrenner, M., see Nuernberg, B. 476
Korf. J., see Wolf, J.H. 496
Kotaki, H., see Ohtani, M. 469
Krenger, W., see Honegger, C.G. 463
Krungkrai, J.
—, Yuthavong, Y. and Webster, H.K.
High-performance liquid chromatographic assay for thymidylate synthase from the human malaria parasite, *Plasmodium falciparum* 51
Kutlar, F.
—, Gu, L.H., Hu, H. and Huisman, T.H.J.
Quantitation of hemoglobins Bart's, H, Portland-I, Portland-II and Constant Spring by anion-exchange high-performance liquid chromatography 265
Labadarios, D.
—, Moodie, I.M., Hough, B.J., Burger, J.A. and Samaai, M.S.
Determination of amino acids in rat muscle by gas chromatography 428
Langemann, H., see Honegger, C.G. 463
Larsson, L., see Sonesson, A. 1
Leroux, F., see Sioufi, A. 81
Lindner, W.
—, Rath, M., Stoschitsky, K. and Uray, G.
Enantioselective drug monitoring of (*R*)- and (*S*)-propranolol in human plasma via derivatization with optically acitve (*R,R*)-O,O-diacetyl tartaric acid anhydride 375
Loschi, L., see Girometta, M.A. 117
Lunte, C.E., see Hendrickson, H.P. 401
McKay, G., see Srinivas, N.R. 61
McNamara, P.J., see Jewell, R.C. 257
Mahuzier, G., see Dessalles, M.C. 215
Man in 't Veld, A.J., see Van der Hoorn, F.A.J. 17
Manning, J.N.
—, Sullivan, G.S. and Davis, P.F.
Reversed-phase liquid chromatography of elastin peptides 41
Manzini, S., see Agostini, O. 331
Markowitz, J.S., see Patrick, K.S. 91
Martin, R.J.
— and Downer, R.G.H.
Microassay for N-acetyltransferase activity using high-performance liquid chromatography with electrochemical detection 287
Mason, W.D., see Gillilan, R.B. 232
Mennes, W.C., see Van Pelt, F.N.A.M. 489
Meyer, M.C., see Patrick, K.S. 91
Midha, K.K., see Srinivas, N.R. 61
Mills, G.A., see Chong, W.K. 147
Mocaer, E., see Tsaconas, C. 313
Moneti, G., see Agostini, O. 331
Moodie, I.M., see Labadarios, D. 428
Morgan, A.K., see Benny, A.G. 173
Motohashi, M., see Yamashita, K. 357
Nakagawa, F., see Ohtani, M. 469
Nakagawa, Y.
— and Waku, K.
Separation of the molecular species of diacyl and alkenylacyl subclasses of the methyl ester of dinitrophenylethanolamine glycerophospholipid by high-performance liquid chromatography 239
Nambara, T., see Shimada, K. 247
Niemeijer, N.R., see Gerding, T.K. 125
Nishitateno, K., see Ohtani, M. 469
Noonan, P.K., see Ruggirello, D.A. 73
Nuernberg, B.
—, Kohlbrenner, M., Faulkner, R. and Furst, D.E.
Rapid quantitation of methotrexate and its metabolites in human serum, urine and bile, using solid-phase extraction and high-performance liquid chromatography 476
Oberkotter, L.V.
Analysis of term human milk concentrations of 3,5,3′-triiodo-L-thyronine by high-performance liquid chromatography and radioimmunoassay: correlation with circulat-

ing serum levels in lactating women 445
Ockelford, P.A., see Benny, A.G. 173
O'Donnell, S.R., see Bryan, L.J. 29
Oe, T., see Shimada, K. 247
Ohtani, M.
—, Shibuya, F., Kotaki, H., Uchino, K., Saitoh, Y., Makagawa. F. and Nishitateno, K.
Quantitative determination of buprenorphine and its active metabolite, norbuprenorphine, in human plasma by gas chromatography–chemical ionization mass spectrometry 469
Oizumi, J., see Hayakawa, K. 161
Owens, J.A., see Smeaton, T.C. 434
Padieu, P., see Tsaconas, C. 313
Patrick, K.S.
—, Markowitz, J.S., Jarvi, E.J., Straughn, A.B. and Meyer, M.C.
Gas chromatographic–mass spectrometric analysis of plasma oxybutynin using a deuterated internal standard 91
Pelt, F.N.A.M. van, see Van Pelt, F.N.A.M. 489
Phillips, L.R., see Brown, N.D. 189
Poon, B.T., see Brown, N.D. 189
Pras, M., see Kaplan, B. 450
Rath, M., see Lindner, W. 375
Robinson, J.S., see Smeaton, T.C. 434
Rohrback, E., see Schwabe, D. 177
Rollema, H., see Den Daas, I. 210
Ruggirello, D.A.
—, Noonan, P.K., Gonzalez, M.A., Ho, P. and Wagner, J.G.
Gas chromatographic determination of gallopamil and norgallopamil in human plasma 73
Rumble, F., see Boutagy, J. 483
Rustum, A.M.
Simple and rapid reversed-phase high-performance liquid chromatographic determination of baclofen in human plasma with ultraviolet detection. Application to a pharmacokinetic study 107
Saitoh, Y., see Ohtani, M. 469
Saller, C.F.
— and Czupryna, M.J.
γ-Aminobutyric acid, glutamate, glycine and taurine analysis using reversed-phase high-performance liquid chromatography and ultraviolet detection of dansyl chloride derivatives 167
Samaai, M.S., see Labadarios, D. 428
Schalekamp, M.A.D.H., see Van der Hoorn, F.A.J. 17
Schmitthenner, H.F.
—, Fedorchuk, M. and Walter, D.J.
Resolution of antihypertensive aryloxypropanolamine enantiomers by reversed-phase chromatography of (–)menthyl chloroformate derivatives 197
Schwabe, D.
—, Rohrbach, E. and Köhl, U.
Determination of *m*-iodobenzylguanidine in serum and urine by high-performance liquid chromatography 177
Schwartz, T.M., see Turner, J.G. 275
Scott, D.O., see Hendrickson, H.P. 401
Shibuya, F., see Ohtani, M. 469
Shimada, K.
—, Oe, T., Tanaka, M. and Nambara, T.
Sensitive ferrocene reagents for derivatization of amines for high-performance liquid chromatography with electrochemical detection 247
Singhvi, S.M., see Everett, D.W. 365
Sioufi, A.
—, Leroux, F. and Bubois, J.P.
Determination of xylometazoline in plasma and urine by gas chromatography using a fused-silica capillary column and an electron-capture detector 81
Smeaton, T.C.
—, Owens, J.A. and Robinson, J.S.
Micro-method for measurement of branched-chain keto acid concentrations in plasma from sheep and man 434
Sonesson, A.
—, Bryn, K., Jantzen, E. and Larsson, L.
Gas chromatographic determination of (phosphorylated) 2-keto-3-deoxyoctonic acid, heptoses and glucosamine in bacterial lipopolysaccharides after treatment with hydrofluoric acid, methanolysis and trifluoroacetylation 1
Spustová, V.
Rapid method for the determination of hippurate in biological fluids by high-performance liquid chromatography 440
Srinivas, N.R.
—, Hubbard, J.W., Hawes, E.M., McKay, G. and Midha, K.K.
Enantioselective gas chromatographic assays with electron-capture detection for methoxyphenamine and its three primary metabolites in human urine 61
Steele, R.D., see Case, G.L. 456
Steinmetz, A.
—, Clavey, V., Vu-Dac, N., Kaffarnik, H. and Fruchart, J.-C.
Purification of human apolipoprotein A-IV by fast protein liquid chromatography 154

Stoschitsky, K., see Lindner, W. 375
Straughn, A.B., see Patrick, K.S. 91
Sugata, R.
—, Iwahashi, H., Ishii, T. and Kido, R.
Separation of polyunsaturated fatty acid radicals by high-performance liquid chromatography with electron spin resonance and ultraviolet detection 9
Sullivan, G.S., see Manning, J.N. 41
Szathmary, S.Cs.
Determination of hydroxypropyl-β-cyclodextrin in plasma and urine by size-exclusion chromatography with post-column complexation 99
Takeda, K., see Hosotsubo, H. 204
Tanaka, M., see Shimada, K. 247
Tepper, P.G., see Den Daas, I. 210
—, see Gerding, T.K. 125
Texier-Maugein, J., see Fourche, J. 142
Tsaconas, C.
—, Padieu, P., d'Athis, P., Mocaer, E. and Bromet, N.
Gas chromatographic–mass spectrometric assessment of the pharmacokinetics of amineptine and its main metabolite in volunteers with liver impairment 313
Turner, J.G.
—, Schwartz, T.M. and Brooks, B.R.
Rapid separation of tritiated thyrotropin-releasing hormone and its catabolic products from mouse and human central nervous system tissues by high-performance liquid chromatography with radioactive flow detection 275
Uchino, K., see Ohtani, M. 469
Uray, G., see Lindner, W. 375
Van de Grampel, V.J.M., see Gerding, T.K. 125
Van der Hoorn, F.A.J.
—, Boomsma, F., Man in 't Veld, A.J. and Schalekamp, M.A.D.H.
Determination of catecholamines in human plasma by high-performance liquid chromatography: comparison beteen a new method with fluorescence detection and an established method with electrochemical detection 17
Van Pelt, F.N.A.M.
—, Mennes, W.C., Hassing, I.G.A.M. and Blaauboer, B.J.
Application of high-performance liquid chromatographic analysis of scoparone and its metabolites in the study of cytochrome P450 differentiation in vitro 489
Veenma-van der Duin, L., see Wolf, J.H. 496
Ventura, P., see Girometta, M.A. 117
Vienne, B., see Dessalles, M.C. 215
Vries, J.B. de, see Den Daas, I. 210
Vu-Dac, N., see Steinmetz, A. 154
Wagner, J.G., see Ruggirello, D.A. 73
Waku, K., see Nakagawa, Y. 239
Walker, V., see Chong, W.K. 147
Walter, D.J., see Schmitthenner, H.F. 197
Warrington, S.J., see Everett, D.W. 365
Weavind, G.P., see Chong, W.K. 147
Webster, H.K., see Krungkrai, J. 51
Weinstein, S.H., see Everett, D.W. 365
Wolf, J.H.
—, Veenma-van der Duin, L. and Korf, J.
Automated analysis procedure for valproic acid in blood, serum and brain dialysate by high-performance liquid chromatography with bromomethylmethoxycoumarin as fluorescent label 496
Woollard, G.A.
Therapeutic drug monitoring of flecainide using high-performance liquid chromatography and comparison with fluorescence polarisation immunoassay 409
Yamashita, K.
—, Motohashi, M. and Yashiki, T.
Sensitive high-performance liquid chromatographic determination of ionic drugs in biological fluids with short-wavelength ultraviolet detection using column switching combined with ion-pair chromatography: application to basic compounds 357
Yashiki, T., see Yamashita, K. 357
Yoshiya, I., see Hosotsubo, H. 204
Yuthavong, Y., see Krungkrai, J. 51
Zeeuw, R.A. de, see Gerding, T.K. 125
Ziemniak, J.A., see Boppana, V.K. 385

Subject Index

N-Acetyltransferase
Microassay for N-acetyltransferase activity using HPLC with electrochemical detection 287

α_1-Acid glycoprotein
α_1-Acid glycoprotein HPLC column (EnantioPAC) as a screening tool for protein binding 257

Adrenaline
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29

Alkenylacyl glycerophosphoethanolamine
Separation of the molecular species of diacyl and alkenylacyl subclasses of the methyl ester of dinitrophenylethanolamine glycerophospholipid by HPLC 239

Amineptine
GC–MS assessment of the pharmacokinetics of amineptine and its main metabolite in volunteers with liver impairment 313

Amines
Sensitive ferrocene reagents for derivatization of amines for HPLC with electrochemical detection 247

Amino acids
γ-Aminobutyric acid, glutamate, glycine and taurine analysis using reversed-phase HPLC and UV detection of dansyl chloride derivatives 167

Amino acids
Determination of amino acids in rat muscle by GC 428

Amino acids
Micro-method for measurement of branched-chain keto acid concentrations in plasma from sheep and man 434

Amino acids, phenylthiohydantion derivatives
Isocratic separation of phenylthiohydantion-amino acids by reversed-phase HPLC 161

γ-Aminobutyric acid
γ-Aminobutyric acid, glutamate, glycine and taurine analysis using reversed-phase HPLC and UV detection of dansyl chloride derivatives 167

Anabolic steroids
Determination of methandrostenolone and its metabolites in equine plasma and urine by coupled-column LC with UV detection and confirmation by tandem MS 341

Angiotensin-converting enzyme inhibitor
Body fluid analysis of a phosphonic acid angiotensin-converting enzyme inhibitor using HPLC and post-column derivatization with *o*-phthalaldehyde 135

Antioxidants
LC determination of common water-soluble antioxidants in biological samples 463

Apolipoprotein A-IV
Purification of human apolipoprotein A-IV by fast protein LC 154

Aryloxypropanolamines
Resolution of antihypertensive aryloxypropanolamine enantiomers by reversed-phase chromatography of (−)-menthyl chloroformate 197

Ascorbic acid
LC determination of common water-soluble antioxidants in biological samples 463

Baclofen
Simple and rapid reversed-phase HPLC determination of baclofen in human plasma with UV detection. Application to a pharmacokinetic study 107

Benzalkonium
HPLC assay of benzalkonium in plasma 221

Benzodiazepines
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295

Branched-chain keto acids
Micro-method for meausrement of branched-chain keto acid concentrations in plasma from sheep and man 434

[^{14}C]Bucromarone
HPLC method for the radiometric determination of [^{14}C]bucromarone in human plasma utilizing non-radiolabeled bucromarone as an internal standard 365

Buprenorphine
Quantitative determination of buprenorphine and its active metabolite, norbuprenorphine, in human plasma by GC–chemical ionization MS 469

Catecholamines
Determination of catecholamines in human plasma by HPLC: comparison between a new method with fluorescence detection and

an established method with electrochemical detection 17

Catecholamines
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29

Catecholamines
Determiantion of *m*-iodobenzylguanidine in serum and urine by HPLC 177

Chloroquine
Identification and determination of a carboxylic acid metabolite of chloroquine in human urine by HPLC 189

3-[(7′-Chloro-4′-quinolinyl)amino]pentanoic acid
Identification and determination of a carboxylic acid metabolite of chloroquine in human urine by HPLC 189

Clobazam
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295

Coagulation factors, vitamin K-dependent
Separation of vitamin K-dependent coagulation factors on sulphated cellulose 173

Cyclodextrins
Determination of hydroxypropyl-β-cyclodextrin in plasma and urine by size-exclusion chromatography with post-column complexation 99

Cysteine
LC determination of common water-soluble antioxidants in biological samples 463

Cytochrome P450
Application of HPLC analysis of scoparone and its metabolites in the study of cytochrome P450 differentiation in vitro 489

Deoxythymidylate
HPLC assay for thymidylate synthase from the human malaria parasite, *Plasmodium falciparum* 51

Deoxyuridylate
HPLC assay for thymidylate synthase from the human malaria parasite, *Plasmodium falciparum* 51

N-Desmethylmethoxyphenamine
Enantioselective GC assays with electron-capture detection for methoxyphenamine and its three primary metabolites in human urine 61

O-Desmethylmethoxyphenamine
Enantioselective GC assays with electron-capture detection for methoxyphenamine and its three primary metabolites in human urine 61

2,4-Diamino-N^{10}-methylpteroic acid
Rapid quantitation of methotrexate and its metabolites in human serum, urine and bile, using solid-phase extraction and HPLC 476

Diazepam
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasme 295

Dinitrophenylethanolamine glycerophospholipids
Separation of the molecular species of diacyl and alkenylacyl subclasses of the methyl ester of dinitrophenylethanolamine glycerophospholipids by HPLC 239

Dopamine
Determination of catecholamines in human plasma by HPLC: comparison between a new method with fluorescence detection and an established method with electrochemical detection 17

Dopamine
Microassay for N-acetyltransferase activity using HPLC with electrochemical detection 287

Dopamine agonist N-0437
Analysis of the dopamine agonist N-0437 in rat serum using reversed-phase HPLC with electrochemical detection 210

Elastin
Reversed-phase LC of elastin peptides 41

Enzymes
HPLC assay for thymidylate synthase from the human malaria parasite, *Plasmodium falciparum* 51

Enzymes
Microassay for N-acetyltransferase activity using HPLC with electrochemical detection 287

17-Epimethandrostenolone
Determination of methandrostenolone and its metabolites in equine plasma and urine by coupled-column LC with UV detection and confirmation by tandem MS 341

Epinephrine
Determination of catecholamines in human plasma by HPLC: comparison between a new method with fluorescence detection and an established method with electrochemical detection 17

Ethanol
Simplified GC analysis of ethanol in blood and tissue 228
Ethanolamine glycerophospholipids
Separation of the molecular species of diacyl and alkenylacyl subclasses of the methyl ester of dinitrophenylethanolamine glycerophospholipid by HPLC 239
1-[2-Ethoxy-2-(3′-pyridyl)ethyl]-4-(2′-methoxyphenyl)piperazine
Determination of the antihypertensive drug 1-[2-ethoxy-2-(3′-pyridyl)ethyl]-4-(2′-methoxyphenyl)piperazine (IP/66) in rat and human plasma by HPLC and isotope dilution MS 331
Factor IX
Separation of vitamin K-dependent coagulation factors in sulphated cellulose 173
Fatty acid radicals
Separation of polyunsaturated fatty acid radicals by HPLC with election spin resonance and UV detection 9
Fatty acids
GC fatty acid determination to differentiate *Nocardia asteroides, Mycobacterium fortuitum* and *Mycobacterium chelonei* 142
Fenoldopam
Simplified procedures for the determination of fenoldopam and its metabolites in human plasma by HPLC with electrochemical detection: comparison of manual and robotic sample preparation methods 385
Ferrocene reagents
Sensitive ferrocene reagents for derivatization of amines for HPLC with electrochemical detection 247
Flecainide
Therapeutic drug monitoring of flecainides using HPLC and comparison with fluorescence polarisation immunoassay 409
Flunitrazepam
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295
5-Fluorocytosine
Rapid and sensitive method for the determination of 5-fluorocytosine in human plasma by ion-pair HPLC 183
2-{3-[4-(4-Fluorophenyl)-1-piperazinyl]propyl}-6,7,8,9-tetrahydro-2*H*-naphtho[2,3-*b*][1,4]oxazin-3(4*H*)-one
Sensitive HPLC determination of ionic drugs in biological fluids with short-wavelength UV detection using column switching combined with ion-pair chromatography: application to basic compounds 357
Folates
Determination of reduced folate derivatives in tissue samples by HPLC with fluorimetric detection 456
Gallopamil
GC determination of gallopamil and norgallopamil in human plasma 73
Glucosamine
GC determination of (phosphorylated) 2-keto-3-deoxyoctonic acid, heptoses and glucosamine in bacterial lipopolysaccharides after treatment with hydrofluoric acid, methanolysis and trifluoroacetylation 1
Glutamate
γ-Aminobutyric acid, glutamate, glycine and taurine analysis using reversed-phase HPLC and UV detection of dansyl chloride derivatives 167
Glycine
γ-Aminobutyric acid, glutamate, glycine and taurine analysis using reversed-phase HPLC and UV detection of dansyl chloride derivatives 167
Haemoglobin Bart's
Quantitation of hemoglobins Bart's, H, Portland-I, Portland-II and Constant Spring by anion-exchange HPLC 265
Haemoglobin Constant Spring
Quantitation of hemoglobins Bart's, H, Portland-I, Portland-II and Constant Spring by anion-exchange HPLC 265
Haemoglobin H
Quantitation of hemoglobins Bart's, H, Portland-I, Portland-II and Constant Spring by anion-exchange HPLC 265
Haemoglobin Portland-I
Quantitation of hemoglobins Bart's, H, Portland-I, Portland-II and Constant Spring by anion-exchange HPLC 265
Haemoglobin Portland-II
Quantitation of hemoglobins Bart's, H, Portland-I, Portland-II and Constant Spring by anion-exchange HPLC 265
Heptoses
GC determination of (phosphorylated) 2-keto-3-deoxyoctonic acid, heptoses and glucosamine in bacterial lipopolysaccarides after treatment with hydrofluoric acid, methanolysis and trifluoroacetylation 1
Hippurate
Rapid method for the determination of hip-

purate in biological fluids by HPLC 440
Hormones
Rapid separation of tritiated thyrotropin-releasing hormone and its catabolic products from mouse and human central nervous system tissues by HPLC with radioactive flow detection 275
Hormones
Analysis of term human milk concentrations of 3,5,3′-triido-L-thyronine by HPLC and radioimmunoassay: correlation with circulating serum levels in lactating women 445
9-Hydroxylamine-1,2,3,4-tetrahydroacridine
Identification of 9-hydroxylamine-1,2,3,4-tetrahydroacridine as a hepatic microsomal metabolite of tacrine by HPLC and electrochemistry 401
5-Hydroxymethoxyphenamine
Enantioselective GC assays with electron-capture detection for methoxyphenamine and its three primary metabolites in human urine 61
7-Hydroxymethotrexate
Rapid quantitation of methotrexate and its metabolites in human serum, urine and bile, using solid-phase extraction and HPLC 476
Hydroxypropyl-β-cyclodextrin
Determination of hydroxypropyl-β-cyclodextrin in plasma and urine by size-exclusion chromatography with post-column complexation 99
m-Iodobenzylguanidine
Determination of *m*-iodobenzylguanidine in serum and uring by HPLC 177
Ionic drugs
Sensitive HPLC determination of ionic drugs in biological fluids with short-wavelength UV detection using column switching combined with ion-pair chromatography: application to basic compounds 357
Isoprenaline
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29
Isoscopoletin
Application of HPLC analysis of scoparone and its metabolites in the study of cytochrome P450 differentiation in vitro 489
Isradipine
Determination of isradipine and the oxidative pyridine metabolite in human plasma by HPLC 483
Keto acids
Micro-method for measurement of branched-chain keto acid concentrations in plasma from sheep and man 434
2-Keto-3-deoxyoctonic acid
GC determination of (phosphorylated) 2-keto-3-deoxyoctonic acid, heptoses and glucosamine in bacterial lipopolysaccharides after treatment with hydrofluoric acid, methanolysis and trifluoroacetylation 1
Lipids
Separation of the molecular species of diacyl and alkenylacyl subclassed of the methyl ester of dinitrophenylethanolamine glycerophospholipid by HPLC 239
Lipoproteins
Purification of human apolipoprotein A-IV by fast protein LC 154
Lorazepam
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295
(−)-Menthyl chloroformate
Resolution of antihypertensive aryloxypropanolamine enantiomers by reversed-phase chromatography of (−)-menthyl chloroformate 197
Metanephrine
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29
Methandrostenolone
Determination of methandrostenolone and tis metabolites in equine plasma and urine by coupled-column LC with UV detection and confirmation by tandem MS 341
Methotrexate
Rapid quantitation of methotrexate and its metabolites in human serum, urine and bile, using solid-phase extraction and HPLC 476
3-Methoxy-4-hydroxymandelic acid
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29
3-Methoxy-4-hydroxyphenylethylene glycol
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29
Methoxyphenamine
Enantioselective GC assays with electron-

capture detection for methoxyphenamine and its three primary metabolites in human urine 61

Methyl-2-(4-diphenylmethyl-1-piperazinyl)ethyl (±)-1,4-dihydro-2,6-dimethyl-4-(*m*-nitrophenyl)-3,5-pyridinedicarboxylate
Sensitive HPLC determination of ion drugs in biological fluids with short-wavelength UV detection using column switching combined with ion-pair chromatography: application to basic compounds 357

3-O-Methylisoprenaline
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29

Midazolam
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295

Mycobacterium chelonei
GC fatty acid determination to differentiate *Nocardia asteroides, Mycobacterium fortuitum* and *Mycobacterium chelonei* 142

Mycobacterium fortuitum
GC fatty acid determination to differentiate *Nocardia asteroides, Mycobacterium fortuitum* and *Mycobacterium chelonei* 142

Nesosteine
HPLC assay of nesosteine in human plasma and urine 117

Nocardia asteroides
GC fatty acid determination to differentiate *Nocardia asteroides, Mycobacterium fortuitum* and *Mycobacterium chelonei* 142

Noradrenaline
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29

Norbuprenorphine
Quantitative determination of buprenorphine and its active metabolite, norbuprenorphine, in human plasma by GC–chemical ionization MS 469

Norepinephrine
Determination of catecholamines in human plasma by HPLC: comparison between a new method with fluorescence detection and an established method with electrochemical detection 17

Norgallopamil
GC determination of gallopamil and norgallopamil in human plasma 73

Normetanephrine
Analysis of the O-methylated metabolites of isoprenaline, adrenaline and noradrenaline in physiological salt solutions by HPLC with electrochemical detection 29

Octopamine
Microassay for N-acetyltransferase activity using HPLC with electrochemical detection 287

Organic acids
HPLC method for the rapid profiling of plasma and urinary organic acids 147

Oxazepam
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295

Oxybutynin
GC–MS analysis of plasma oxybutynin using a deuterated internal standard 91

Peptides
Reversed-phase LC of elastin peptides 41

Peptides
Separation of peptides from sodium dodecyl sulphate by HPLC 450

Perphenazine
Determination of perphenazine in serum by LC 215

Phosphonic acid angiotensin-convering enzyme inhibitor
Body fluid analysis of a phosphonic acid angiotensin-converting enzyme inhibitor using HPLC and post-column derivatization with *o*-phthalaldehyde 135

Piroxicam
Rapid analysis of piroxicam in dog, rat and human plasma by HPLC 232

Plasmalogen
Separation of the molecular species of diacyl and alkenylacyl subclasses of the methyl ester of dinitrophenylethanolamine glycerophospholipid by HPLC 239

Polyunsaturated fatty acids
Separation of polyunsaturated fatty acid radicals by HPLC with electron spin resonance and UV detection 9

Propranolol
Enantioselective drug monitoring of (*R*)- and (*S*)-propranolol in hyman plasma via derivatization with optically active (*R,R*)-O,O-diacetyl tartaric acid anhydride 375

2-(N-Propyl-N-2-thienylethylamino)-5-

hydroxytetralin
Determination of enantiomeric purity of the new D-2 dopamine agonist 2-(N-propyl-N-2-thienylethylamino)-5-hydroxytetralin (N-0437) by reversed-phase HPLC after pre-column derivatization with D(+)-glucuronic acid 125
2-(N-Propyl-N-2-thienylethylamino)-5-hydroxytetralin
Analysis of the dopamine agonist N-0437 in rat serum using reversed-phase HPLC with electrochemical detection 210
Proteins
α-Acid glycoprotein HPLC column (EnantioPAC) as a screening tool for protein binding 257
Pyridines
Determination of isradipine and the oxidative pyridine metabolite in human plasma by HPLC 483
Pyrimidines
Rapid and sensitive method for the determination of 5-fluorocytosine in human plasma by ion-pair HPLC 183
Reduced glutathione
LC determination of common water-soluble antioxidants in biological samples 463
Scoparone
Application of HPLC analysis of scoparone and its metabolites in the study of cytochrome P450 differentiation in vitro 489
Scopoletin
Application of HPLC analysis of scoparone and its metabolites in the study of cytochrome P450 differentiation in vitro 489
Serotonin
Microassay for N-acetyltransferase activity using HPLC with electrochemical detection 287
Sodium dodecyl sulphate
Separation of peptides from sodium dodecyl sulphate by HPLC 450
Steroids
Determination of methandrostenolone and its metabolites in equine plasma and urine by coupled-column LC with UV detection and confirmation by tandem MS 341
N-Succinimidyl 3-ferrocenylpropionate
Sensitive ferrocene reagents for derivatization of amines for HPLC with electrochemical detection 247
Sugars
GC determination of (phosphorylated) 2-keto-3-deoxyoctonic acid, heptoses and glucosamine in bacterial lipopolysaccharides after treatment with hydrofluoric acid, methanolysis and trifluoroacetylation 1
Sulphated cellulose
Separation of vitamin K-dependent coagulation factors on sulphated cellulose 173
Tacrine
Identification of 9-hydroxylamine-1,2,3,4,-tetrahydroacridine as a hepatic microsomal metabolite of tacrine by HPLC and electrochemistry 401
Taurine
γ-Aminobutyric acid, glutamate, glycine and taurine analysis using reversed-phase HPLC and UV detection of dansyl chloride derivatives 167
Thiamylal
Measurement of thiamylal in human plasma using reversed-phase HPLC 204
Thymidylate synthase
HPLC assay for thymidylate synthase from the human malaria parasite, *Plasmodium falciparum* 51
Thyroid hormones
Analysis of term human milk concentrations of 3,5,3′-triido-L-thyronine by HPLC and radioimmunoassay: correlation with circulating serum levels in lactating women 445
Thyrotropin-releasing hormone
Rapid separation of tritiated thyrotropin-releasing hormone and its catabolic products from mouse and human central nervous system tissues by HPLC with radioactive flow detection 275
Triazolam
Capillary GC–MS method for the identification and quantification of some benzodiazepines and their unconjugated metabolites in plasma 295
3,5,3′-Triido-L-thyronine
Analysis of term human milk concentrations of 3,5,3′-triido-L-thyronine by HPLC and radioimmunoassay: correlation with circulating serum levels in lactating women 445
Uric acid
LC determination of common water-soluble antioxidants in biological samples 463
Valproic acid
Automated analysis procudure for valproic acid in blood, serum and brain dialysate by HPLC with bromomethylmethoxy-coumarin as fluorescent label 496

Vancomycin
Rapid and specific method for the determination of vancomycin in plasma by HPLC on an aminopropyl column 421
Xylometazoline
Determination of xylometazoline in plasma and urine by GC using a fused-silica capillary column and an electron-capture detector 81